

LIPSCHITZNESS OF *-HOMOMORPHISMS BETWEEN C*-METRIC ALGEBRAS

WEI WU

ABSTRACT. A C^* -metric algebra consists of a unital C^* -algebra and a Leibniz Lip-norm on the C^* -algebra. We show that if the Lip-norms concerned are lower semicontinuous, then any unital *-homomorphism from a C^* -metric algebra to another one is necessarily Lipschitz. It results that the free product of two Lipschitz unital *-homomorphisms between C^* -metric algebras coming from *-filtrations is still a Lipschitz unital *-homomorphism.

1. INTRODUCTION

Originated in Kantorovič's work [12, 13] and Connes' observation [4], compact quantum metric space was introduced by Rieffel as the noncommutative analogue of compact metric space [24, 25]. It consists of an order unit space and a Lip-norm on it, where the Lip-norm plays the role of the usual Lipschitz seminorms for ordinary compact metric spaces [26, 22, 29]. From this, a way to view "matrix algebras converge to the sphere" is established by defining the Gromov-Hausdorff distance between compact quantum metric spaces [27, 28]. See [14, 18, 39, 16, 19, 31] for further discussion.

To precisely formulate the statements in the literature of high-energy physics and string-theory, such as "here are the vector bundles over the matrix algebras that correspond to the monopole bundles over the sphere" in [32, 2, 10] etc., Rieffel found that algebraic conditions on the Lip-norm, such as the Leibniz rule, play a crucial role [30, 31].

At the investigation of the structure of unital C^* -algebras with complexified Lip-norms, Kerr proposed a notion of dimension for these C^* -algebras, along with two dynamical entropies. In this environment, the Leibniz rule on the Lip-norm is also of central importance [15].

So unital C^* -algebras with Leibniz Lip-norms form an important class of compact quantum metric spaces. For these class of objects, it is essential that the maps preserve the given algebraic structure and seminorms in same way. The morphisms between two such objects were discussed in [15]: Lipschitz unital *-homomorphisms. It is natural to ask what conditions are needed to guarantee that a unital *-homomorphism from one such object to another is Lipschitz. In this note we show

2000 *Mathematics Subject Classification.* Primary 46L87; Secondary 46L05.

Key words and phrases. C^* -metric algebra, unital *-homomorphism, lower semicontinuous seminorm, Leibniz seminorm, reduced free product, Lipschitz map.

The research was supported by the Shanghai Leading Academic Discipline Project (Project No. B407), and the National Natural Science Foundation of China (Grant No. 10671068).

Theorem 1.1. *Let (A_1, L_1) and (A_2, L_2) be two C^* -metric algebras with lower semicontinuous Lip-norms. If T is a unital $*$ -homomorphism from (A_1, L_1) into (A_2, L_2) , then T is Lipschitz.*

(Precise definitions are given in Section 2.)

In fact, any Lip-norm on a unital C^* -algebra may be replaced by a lower semicontinuous Lip-norm according to theorem 4.1 in [25]. Therefore, unital C^* -algebras with lower semicontinuous Lip-norms are a class of C^* -algebras that deserve to be studied further.

We recall in Section 2 some terminology and some typical constructions of unital C^* -algebras with lower semicontinuous Leibniz Lip-norms. We prove Theorem 1.1 in Section 3, and as an application we show, in Section 4, that the free product of two Lipschitz unital $*$ -homomorphisms between C^* -metric algebras coming from $*$ -filtrations is still a Lipschitz unital $*$ -homomorphism.

Acknowledgements. I am grateful to Hanfeng Li for comments.

2. C^* -METRIC ALGEBRAS

We recall [31] that a *Lip-norm* on a unital C^* -algebra A is a seminorm L on A , possibly taking the value $+\infty$, such that

- (1) $L(1_A) = 0$, where 1_A is the identity of A ,
- (2) $L(a^*) = L(a)$ for all $a \in A$,
- (3) the metric ρ_L on $S(A)$ given by

$$\rho_L(\varphi, \psi) = \sup\{|\varphi(a) - \psi(a)| : L(a) \leq 1\}$$

induces the weak*-topology on $S(A)$.

When L is a Lip-norm on a unital C^* -algebra A , the metric space $(S(A), \rho_L)$ is compact. So the diameter $\text{diam}(A, L)$ of (A, L) , given by the diameter of $S(A)$ with respect to ρ_L , is finite. If $L(a) = 0$ and the condition 3 above holds, then $L(na) = 0$ for any $n \in \mathbb{N}$, and so $|\varphi(a) - \psi(a)| \leq \frac{1}{n}\rho_L(\varphi, \psi)$ for all $\varphi, \psi \in S(A)$ and $n \in \mathbb{N}$. Hence $\varphi(a) = \psi(a)$ for all $\varphi, \psi \in S(A)$. Therefore, $a \in \mathbb{C}1_A$. Let $\varphi, \psi \in S(A)$ with $\varphi \neq \psi$. Since the weak*-topology on $S(A)$ is Hausdorff, $\rho_L(\varphi, \psi) > 0$ if the metric ρ_L gives $S(A)$ the weak*-topology. So there is an $a \in A$ with $L(a) \leq 1$ such that $\varphi(a) \neq \psi(a)$. This indicates that L separates the state space $S(A)$ of A .

A seminorm L on a unital C^* -algebra A is said to be *lower semicontinuous* if for one $r \in \mathbb{R}_{>0}$, hence for all $r > 0$, the set $\{a \in A : L(a) \leq r\}$ is norm-closed in A . Given a seminorm on a unital C^* -algebra A , we denote the set $\{a \in A : L(a) < \infty\}$ of Lipschitz elements in A by $\mathcal{A}$. The seminorm L is said to be *Leibniz* if it satisfies the Leibniz rule

$$L(ab) \leq L(a)\|b\| + \|a\|L(b)$$

for all $a, b \in \mathcal{A}$.

Now we define

Definition 2.1. A *C^* -metric algebra* is a pair (A, L) consisting of a unital C^* -algebra A and a Leibniz Lip-norm L on A .

Basically, compact metric spaces provide the prototype of C^* -metric algebras.

Example 2.2. Let (X, ρ) be an ordinary compact metric space, and let L_ρ denote the Lipschitz seminorm on $C(X)$. Then $(C(X), L_\rho)$ is a C^* -metric algebra and L_ρ is lower semicontinuous.

One important class of examples of C^* -metric algebras comes from ergodic actions of a compact group on a unital C^* -algebra [24, 31].

Example 2.3. Let A be a unital C^* -algebra, let G be a compact group, and let α be a strongly continuous ergodic action of G on A by automorphisms. We assume that G is equipped with a length function ℓ . Then the seminorm

$$L(a) = \sup \{ \|\alpha_x(a) - a\| / \ell(x) : x \in G \setminus \{e\} \}, \quad a \in A,$$

is a lower semicontinuous Leibniz Lip-norm on A .

In [20], a generalization to ergodic actions of co-amenable compact quantum group was given. By viewing an ergodic action of a compact group on a unital C^* -algebra as the translation action of the group on a noncommutative homogeneous space of it, Hanfeng Li recently extended the construction above to the locally compact groups which satisfy certain conditions [21].

Example 2.4. Given a spectral triple (A, H, D) , where (H, D) is an unbounded Fredholm module over a unital C^* -algebra A [5, 6], we can define a seminorm, L , on A by

$$L(a) = \|[D, a]\|.$$

Let σ be a state of A . Then L is a Lip-norm if and only the set $\{a \in A : L(a) \leq 1, \sigma(a) = 0\}$ is a norm-totally-bounded subset of A [22]. In this case, (A, L) is a C^* -metric algebra with lower semicontinuous Lip-norm.

In fact, every C^* -metric algebra can be obtained from the Dirac operator approach [29]. Finally, we make:

Definition 2.5. Let (A_1, L_1) and (A_2, L_2) be two C^* -metric algebras. By a **-homomorphism from (A_1, L_1) to (A_2, L_2)* we mean a *-homomorphism T from C^* -algebra A_1 to C^* -algebra A_2 such that $T(\mathcal{A}_1) \subseteq \mathcal{A}_2$. A map $\alpha : A_1 \rightarrow A_2$ is said to be *Lipschitz* if there is a constant $\mu \geq 0$ such that

$$L_2(\alpha(a)) \leq \mu L_1(a)$$

for all $a \in A_1$.

3. *-HOMOMORPHISMS BETWEEN C*-METRIC ALGEBRAS

Let A be a unital C^* -algebra with a *Lipschitz seminorm* L , a seminorm which may take value $+\infty$, has its null-space the scalar multiples of the identity, and satisfies $L(a^*) = L(a)$ for all $a \in A$. We can define a metric ρ_L on the state space $S(A)$ of A by

$$\rho_L(\varphi, \psi) = \sup \{ |\varphi(a) - \psi(a)| : L(a) \leq 1 \}, \quad \varphi, \psi \in S(A).$$

Usually this metric may take value $+\infty$. From this metric we can define a seminorm L_{ρ_L} on A by the formula

$$L_{\rho_L}(a) = \sup \left\{ \frac{|\varphi(a) - \psi(a)|}{\rho_L(\varphi, \psi)} : \varphi \neq \psi, \varphi, \psi \in S(A) \right\}, \quad a \in A.$$

This is still a Lipschitz seminorm.

The dual Banach space to $A/(\mathbb{C}1_A)$ for the quotient norm $\|\cdot\|^\sim$ is just the subspace A'^0 of the dual space A' consisting of those $f \in A'$ such that $f(1_A) = 0$. Similar to [25], we define

$$L'(f) = \sup \{ |f(a)| : L(a) \leq 1, a \in A \}, \quad f \in A'.$$

By proposition 3.7 of [23] and proposition 4.2 of [38], for any $g \in A'^0$ with $\|g\| \leq 2$ there are $\varphi_1, \varphi_2, \varphi_3, \varphi_4 \in S(A)$ such that $g = (\varphi_1 - \varphi_2) + i(\varphi_3 - \varphi_4)$. So

$$\begin{aligned} L'(g) &= \sup\{|g(a)| : L(a) \leq 1, a \in A\} \\ &= \sup\{|(\varphi_1(a) - \varphi_2(a)) + i(\varphi_3(a) - \varphi_4(a))| : L(a) \leq 1, a \in A\} \\ &\leq \sup\{|\varphi_1(a) - \varphi_2(a)| : L(a) \leq 1, a \in A\} \\ &\quad + \sup\{|\varphi_3(a) - \varphi_4(a)| : L(a) \leq 1, a \in A\} \\ &= \rho_L(\varphi_1, \varphi_2) + \rho_L(\varphi_3, \varphi_4), \end{aligned}$$

and

$$\begin{aligned} \rho_L(\varphi_1, \varphi_2) + \rho_L(\varphi_3, \varphi_4) &= \sup\{|\varphi_1(a) - \varphi_2(a)| : L(a) \leq 1, a \in A\} \\ &\quad + \sup\{|\varphi_3(a) - \varphi_4(a)| : L(a) \leq 1, a \in A\} \\ &= \sup\{|g(a) + g^*(a)|/2 : L(a) \leq 1, a \in A\} \\ &\quad + \sup\{|g(a) - g^*(a)|/2 : L(a) \leq 1, a \in A\} \\ &\leq 2\sup\{|g(a)| : L(a) \leq 1, a \in A\} \\ &= 2L'(g). \end{aligned}$$

Denote

$$\mathcal{L}_1 = \{a \in A : L(a) \leq 1\}.$$

Then $\mathcal{L}_1$ is convex and balanced, and the bipolar theorem says that $\overline{\mathcal{L}_1}^{\|\cdot\|} = \mathcal{L}_1^{\circ\circ}$. Now suppose that L is lower semicontinuous. Then we get

$$\mathcal{L}_1 = \mathcal{L}_1^{\circ\circ}.$$

If $L_{\rho_L}(a) \leq 1$, then $|\varphi(a) - \psi(a)| \leq \rho_L(\varphi, \psi)$ for all $\varphi, \psi \in S(A)$. So $|g(a)| \leq 2L'(g)$ for all $g \in A'^0$ with $\|g\| \leq 2$, and this also implies that $|g(a)| \leq 2L'(g)$ for all $g \in A'^0$. Thus $|g(a/2)| \leq 1$ for all $g \in A'$ with $L'(g) \leq 1$. By definition, we obtain that $a/2 \in \mathcal{L}_1^{\circ\circ} = \mathcal{L}_1$. So

$$L(a) \leq 2L_{\rho_L}(a).$$

If $L(a) \leq 1$, then $a \in \mathcal{L}_1^{\circ\circ}$. So $|f(a)| \leq 1$ for all $f \in A'$ with $L'(f) \leq 1$, and hence $|f(a)| \leq L'(f)$ for all $f \in A'^0$. In particular, we have that $|\varphi(a) - \psi(a)| \leq L'(\varphi - \psi) = \rho_L(\varphi, \psi)$ for $\varphi, \psi \in S(A)$. Thus $L_{\rho_L}(a) \leq 1$. So

$$L_{\rho_L}(a) \leq L(a).$$

Therefore, we have

Lemma 3.1. *Let A be a unital C^* -algebra with a lower semicontinuous Lipschitz seminorm L . Then*

$$L_{\rho_L}(a) \leq L(a) \leq 2L_{\rho_L}(a),$$

for all $a \in A$.

Suppose A_1 and A_2 are two unital C^* -algebras with C^* -norms $\|\cdot\|_1$ and $\|\cdot\|_2$, respectively. Let L_1 and L_2 be Lipschitz seminorms on A_1 and A_2 , respectively, which are Leibniz. On $\mathcal{A}_1$ and $\mathcal{A}_2$, we can define new norms by

$$M_1(a) = \|a\|_1 + L_1(a), \quad a \in \mathcal{A}_1,$$

$$M_2(b) = \|b\|_2 + L_2(b), \quad b \in \mathcal{A}_2,$$

respectively. Then $(\mathcal{A}_1, M_1)$ and $(\mathcal{A}_2, M_2)$ are normed algebras since L_1 and L_2 are Leibniz. We denote by $\mathfrak{A}_1$ and $\mathfrak{A}_2$ the completions of $\mathcal{A}_1$ and $\mathcal{A}_2$ with respect to the norms M_1 and M_2 , respectively.

Let $\{a_n\}$ be a sequence in $\mathcal{A}_1$ such that $\sum_{n=1}^{\infty} M_1(a_n)$ converges. Then both $\sum_{n=1}^{\infty} \|a_n\|_1$ and $\sum_{n=1}^{\infty} L_1(a_n)$ are convergent. Since $\mathcal{A}_1$ is complete in the norm $\|\cdot\|_1$, there is an $a \in \mathcal{A}_1$ such that $a = \sum_{n=1}^{\infty} a_n$ in the norm $\|\cdot\|_1$. For any $k \in \mathbb{N}$, we have

$$L_1\left(\sum_{n=1}^k a_n\right) \leq \sum_{n=1}^k L_1(a_n) \leq \sum_{n=1}^{\infty} L_1(a_n).$$

Now assume that L_1 is lower semicontinuous. Then we have

$$L_1(a) \leq \sum_{n=1}^{\infty} L_1(a_n) < \infty.$$

Thus $a \in \mathcal{A}_1$. Similarly, we have that $L_1\left(\sum_{n=k+1}^{\infty} a_n\right) \leq \sum_{n=k+1}^{\infty} L_1(a_n)$ for $k \in \mathbb{N}$. So

$$\begin{aligned} M_1\left(a - \sum_{n=1}^k a_n\right) &= \left\|a - \sum_{n=1}^k a_n\right\|_1 + L_1\left(a - \sum_{n=1}^k a_n\right) \\ &\leq \sum_{n=k+1}^{\infty} \|a_n\|_1 + \sum_{n=k+1}^{\infty} L_1(a_n). \end{aligned}$$

Thus $\lim_{k \rightarrow \infty} M_1\left(a - \sum_{n=1}^k a_n\right) = 0$, i.e., $\sum_{n=1}^{\infty} a_n$ converges to a in the norm M_1 . So $\mathcal{A}_1$ is complete in the norm M_1 . Therefore, $\mathfrak{A}_1 = \mathcal{A}_1$. Similarly, we have that $\mathfrak{A}_2 = \mathcal{A}_2$ if L_2 is lower semicontinuous. Moreover, in this case we have

Proposition 3.2. *If α is a *-homomorphism from C*-algebra A_1 into C*-algebra A_2 such that $\alpha(\mathcal{A}_1) \subseteq \mathcal{A}_2$, then there is a constant $\lambda > 0$ such that*

$$M_2(\alpha(a)) \leq \lambda M_1(a)$$

for all $a \in \mathcal{A}_1$.

Proof. Since both the norms $\|\cdot\|_1$ and $\|\cdot\|_2$ and the seminorms L_1 and L_2 have an isometric involution, the involutions on $\mathcal{A}_1$ and $\mathcal{A}_2$ are isometric with the norms M_1 and M_2 , respectively. And so $(\mathcal{A}_1, M_1)$ and $(\mathcal{A}_2, M_2)$ are Banach *-algebras with isometric involutions. That α is a *-homomorphism implies that $\alpha(\mathcal{A}_1)$ is a *-subalgebra of $\mathcal{A}_2$. Also from the relation $\mathcal{A}_2 \subseteq A_2$, we get the restriction of the C*-norms on A_2 to $\mathcal{A}_2$, and it makes $\mathcal{A}_2$ into a pre-C*-algebra [37]. This shows that $(\mathcal{A}_2, M_2)$ is an A*-algebra. By theorem 23.11 in [8], α is continuous as a map from $(\mathcal{A}_1, M_1)$ into $(\mathcal{A}_2, M_2)$. Thus, there is a constant $\lambda > 0$ such that $M_2(\alpha(a)) \leq \lambda M_1(a)$ for all $a \in \mathcal{A}_1$. $\square$

Assume that T is a unital positive linear map from C*-algebra A_1 into C*-algebra A_2 . Let $S(A_1)$ and $S(A_2)$ denote the state spaces of A_1 and A_2 , respectively. For any $\varphi \in S(A_2)$, we define

$$\sigma_T(\varphi)(a) = \varphi(Ta), \quad a \in A_1.$$

Then σ_T is an affine map from $S(A_2)$ into $S(A_1)$. Moreover, if T maps A_1 onto A_2 , then σ_T is injective; if σ_T is a surjection, then T is an injection.

Lemma 3.3. *Let (A_1, L_1) and (A_2, L_2) be two C^* -metric algebras with lower semicontinuous Lip-norms. If T is a unital $*$ -homomorphism from (A_1, L_1) into (A_2, L_2) , then there is a Lipschitz affine map σ from metric space $(S(A_2), \rho_{L_2})$ into metric space $(S(A_1), \rho_{L_1})$ such that*

$$\varphi(Ta) = \sigma(\varphi)(a), \quad a \in A_1, \varphi \in S(A_2).$$

Proof. Suppose T is a unital $*$ -homomorphism from A_1 into A_2 . Set $\sigma = \sigma_T$. For any $\varphi_0 \in S(A_1)$ and $b \in A_1$ with $L_1(b) \leq 1$, define

$$f_{\varphi_0, b}(\varphi) = \varphi(b) - \varphi_0(b), \quad \varphi \in S(A_1).$$

Clearly $f_{\varphi_0, b}$ is a continuous affine function on $S(A_1)$. To each $a \in A_1$, we get a function $\hat{a}$ on $S(A_1)$ defined by $\hat{a}(\varphi) = \varphi(a)$. By the basic representation theorem of Kadison [11] (see theorem II.1.8 of [1]), there is a self-adjoint $a_{\varphi_0, b} \in A_1$ such that $f_{\varphi_0, b} = \hat{a}_{\varphi_0, b}$ for every $b = b^* \in A_1$.

For any $\varphi \in S(A_1)$ and $b = b^* \in A_1$ with $L_1(b) \leq 1$, we have

$$|\hat{a}_{\varphi_0, b}(\varphi)| = |\varphi(b) - \varphi_0(b)| \leq \rho_{L_1}(\varphi, \varphi_0) \leq \text{diam}(A_1, L_1).$$

So $\|a_{\varphi_0, b}\|_1 \leq \text{diam}(A_1, L_1)$ by proposition II.1.7 of [1]. And since L_1 is lower semicontinuous, we get

$$\begin{aligned} L_1(a_{\varphi_0, b}) &\leq 2L_{\rho_{L_1}}(a_{\varphi_0, b}) \\ &= 2\sup \left\{ \frac{|\varphi_1(a_{\varphi_0, b}) - \varphi_2(a_{\varphi_0, b})|}{\rho_{L_1}(\varphi_1, \varphi_2)} : \varphi_1 \neq \varphi_2, \varphi_1, \varphi_2 \in S(A_1) \right\} \\ &= 2\sup \left\{ \frac{|\hat{a}_{\varphi_0, b}(\varphi_1) - \hat{a}_{\varphi_0, b}(\varphi_2)|}{\rho_{L_1}(\varphi_1, \varphi_2)} : \varphi_1 \neq \varphi_2, \varphi_1, \varphi_2 \in S(A_1) \right\} \\ &= 2\sup \left\{ \frac{|\varphi_1(b) - \varphi_2(b)|}{\rho_{L_1}(\varphi_1, \varphi_2)} : \varphi_1 \neq \varphi_2, \varphi_1, \varphi_2 \in S(A_1) \right\} \\ &= 2L_{\rho_{L_1}}(b) \leq 2L_1(b) \leq 2 \end{aligned}$$

by Lemma 3.1. Thus

$$M_1(a_{\varphi_0, b}) \leq \text{diam}(A_1, L_1) + 2.$$

So the set $\{a_{\varphi_0, b} : b = b^* \in A_1, L_1(b) \leq 1\}$ is bounded in A_1 with respect to the norm M_1 . By Proposition 3.2, the set $\{Ta_{\varphi_0, b} : b = b^* \in A_1, L_1(b) \leq 1\}$ is bounded in A_2 under the norm M_2 . Then there is a constant $K > 0$ such that $M_2(Ta_{\varphi_0, b}) \leq K$ for all $b = b^* \in A_1$ with $L_1(b) \leq 1$.

Now for any $\psi_1, \psi_2 \in S(A_2)$ and any $b = b_1 + ib_2 \in A_1$ with $L_1(b) \leq 1$ and $b_i = b_i^* \in A_1$ for $i \in \{1, 2\}$, we have that $L_1(b_1) \leq 1$ and $L_1(b_2) \leq 1$, and so

$$\begin{aligned} |\sigma(\psi_1)(b_i) - \sigma(\psi_2)(b_i)| &= |[\sigma(\psi_1)(b_i) - \varphi_0(b_i)] - [\sigma(\psi_2)(b_i) - \varphi_0(b_i)]| \\ &= |\hat{a}_{\varphi_0, b_i}(\sigma(\psi_1)) - \hat{a}_{\varphi_0, b_i}(\sigma(\psi_2))| \\ &= |\hat{T}a_{\varphi_0, b_i}(\psi_1) - \hat{T}a_{\varphi_0, b_i}(\psi_2)| \\ &\leq L_{\rho_{L_2}}(Ta_{\varphi_0, b_i})\rho_{L_2}(\psi_1, \psi_2) \\ &\leq L_2(Ta_{\varphi_0, b_i})\rho_{L_2}(\psi_1, \psi_2) \\ &\leq M_2(Ta_{\varphi_0, b_i})\rho_{L_2}(\psi_1, \psi_2) \\ &\leq K\rho_{L_2}(\psi_1, \psi_2), \end{aligned}$$

by Lemma 3.1. From this we obtain

$$\begin{aligned} |\sigma(\psi_1)(b) - \sigma(\psi_2)(b)| &\leq |\sigma(\psi_1)(b_1) - \sigma(\psi_2)(b_1)| + |\sigma(\psi_1)(b_2) - \sigma(\psi_2)(b_2)| \\ &\leq 2K\rho_{L_2}(\psi_1, \psi_2). \end{aligned}$$

Therefore, $\rho_{L_1}(\sigma(\psi_1), \sigma(\psi_2)) \leq 2K\rho_{L_2}(\psi_1, \psi_2)$. Hence σ is Lipschitz. $\square$

We are ready to prove Theorem 1.1.

Proof of Theorem 1.1 By Lemma 3.3, there exists an affine map σ from metric space $(S(A_2), \rho_{L_2})$ into metric space $(S(A_1), \rho_{L_1})$ such that

$$\varphi(Ta) = \sigma(\varphi)(a), \quad a \in A_1, \varphi \in S(A_2),$$

and

$$\rho_{L_1}(\sigma(\psi_1), \sigma(\psi_2)) \leq K\rho_{L_2}(\psi_1, \psi_2), \quad \psi_1, \psi_2 \in S(A_2),$$

for some positive constant K . So for any $a \in A_1$, we have

$$\begin{aligned} L_2(Ta) &\leq 2L_{\rho_{L_2}}(Ta) \\ &= 2\sup \left\{ \frac{|\varphi(Ta) - \psi(Ta)|}{\rho_{L_2}(\varphi, \psi)} : \varphi \neq \psi, \varphi, \psi \in S(A_2) \right\} \\ &= 2\sup \left\{ \frac{|\sigma(\varphi)(a) - \sigma(\psi)(a)|}{\rho_{L_2}(\varphi, \psi)} : \varphi \neq \psi, \varphi, \psi \in S(A_2) \right\} \\ &= 2\sup \left\{ \frac{|\sigma(\varphi)(a) - \sigma(\psi)(a)|}{\rho_{L_1}(\sigma(\varphi), \sigma(\psi))} \cdot \frac{\rho_{L_1}(\sigma(\varphi), \sigma(\psi))}{\rho_{L_2}(\varphi, \psi)} : \varphi \neq \psi, \varphi, \psi \in S(A_2) \right\} \\ &\leq 2KL_1(a) \end{aligned}$$

by Lemma 3.1. Therefore, T is Lipschitz. $\square$

4. AN APPLICATION

Let A be a unital C^* -algebra. A $*$ -filtration $\{A_n\}$ of A is a sequence of finite-dimensional subspaces which satisfy

- (1) $A_0 = \mathbb{C}1_A$, $A_n^* = A_n$,
- (2) $A_m \subset A_n$ if $m < n$,
- (3) $A_m A_n \subseteq A_{m+n}$,
- (4) $A = \overline{\cup_{n=0}^{\infty} A_n}$

[35]. Given a faithful state σ on A . Let (π, H, ξ) be the faithful GNS representation of (A, σ) . We identify A with the corresponding linear space of vectors in H . We let $\|\cdot\|$ denote the operator norm of $\pi(A)$, and $\|\cdot\|_2$ denote the vector norm of A . Viewing each A_n as a finite-dimensional subspace of H , we obtain a filtration of H in the sense that $\{A_n\}$ is an increasing sequence of finite-dimensional subspaces of H and $\cup_{n=0}^{\infty} A_n$ is dense in H [35]. Let Q_n denote the orthogonal projection of H onto A_n . Set $P_n = Q_n - Q_{n-1}$ for $n \geq 1$, and $P_0 = Q_0$. We define

$$D = \sum_{n=1}^{\infty} nP_n.$$

Then D is a unbounded linear operator on H with domain $\mathcal{A} = \cup_{n=0}^{\infty} A_n$. The linear functional $f(h) = \langle Dh, k \rangle$ on $\mathcal{A}$ is bounded if and only if $k \in \mathcal{A}$. It is also clear that $\langle Dh, k \rangle = \langle h, Dk \rangle$ for all $h, k \in \mathcal{A}$. So D is self-adjoint [7]. By lemma 1.1 of [22], $\mathcal{A} = \{a \in A : [D, \pi(a)] \text{ is bounded}\}$. Furthermore D has a finite-dimensional kernel, and the inverse D^{-1} (defined on the orthogonal complement of D 's kernel)

is compact. So $(\mathcal{A}, H, D)$ is a spectral triple [4, 5, 6, 17, 33]. Using this spectral triple, we can define a seminorm L on A , possible taking the value $+\infty$, by

$$L(a) = \|[D, \pi(a)]\|, \quad a \in A.$$

For all $a, b \in A$, we have

$$\begin{aligned} L(ab) &= \|[D, \pi(ab)]\| = \|[D, \pi(a)]\pi(b) + \pi(a)[D, \pi(b)]\| \\ &\leq \|[D, \pi(a)]\|\|\pi(b)\| + \|\pi(a)\|\|[D, \pi(b)]\| \\ &= L(a)\|b\| + \|a\|L(b). \end{aligned}$$

So L is Leibniz. By main theorem 1.2 in [22], we have

Proposition 4.1. *If there is a constant C such that*

$$\|P_m \pi(P_k(a)) P_n\| \leq C \|P_k(a)\|_2$$

for all $a \in \mathcal{A}$ and $m, n, k \in \mathbb{N}$, then (A, L) is a C^ -metric algebra with lower semicontinuous Lip-norm.*

A condition of this kind is called a *Haagerup-type condition* with constant C . In this case, we will call (A, L) the *C^* -metric algebra coming from $(\{A_n\}, \sigma)$* (with constant C).

Suppose that (A^1, L_1) and (A^2, L_2) are two C^* -metric algebras coming from $(\{A_n^1\}, \sigma_1)$ and $(\{A_n^2\}, \sigma_2)$, respectively, and both with constant C . Let

$$(A, \sigma) = (A^1, \sigma_1) * (A^2, \sigma_2)$$

be the reduced free-product C^* -algebra with the faithful state σ [34, 36, 3]. We define a $*$ -filtration $\{A_n\}$ on (A, σ) by setting A_n to be the linear span of all products $A_{n_1}^{\iota_1} \cdots A_{n_k}^{\iota_k}$ with each $\iota_j \in \{1, 2\}$, with $\iota_j \neq \iota_{j+1}$ for $1 \leq j \leq k-1$, and with $\sum_{j=1}^k n_j \leq n$. We let $(\pi_\iota, H^\iota, \xi_\iota)$ denote the faithful GNS representation of (A^ι, σ_ι) for $\iota \in \{1, 2\}$, and we let (π, H, ξ) denote the faithful GNS representation of (A, σ) . We let $\{P_n\}$ be the family of mutually orthogonal projections corresponding to the filtration $\{A_n\}$ as above. Let D be the Dirac operators coming from $(A, \{A_n\}, \sigma)$. Then

$$D = \sum_{n=1}^{\infty} n P_n,$$

and the corresponding lower semicontinuous seminorm on A is

$$L(a) = \|[D, \pi(a)]\|, \quad a \in A.$$

By theorem 6.1 of [22], we obtain

Proposition 4.2. *If σ_1 and σ_2 are traces, then (A, L) is a C^* -metric algebra coming from $(\{A_n\}, \sigma)$ with constant $\sqrt{5}C$, the reduced C^* -norm on A is the operator norm for the GNS representation for σ on H , and $(H, \xi) = (H^1, \xi_1) * (H^2, \xi_2)$.*

From the discussion above, we see that $\mathcal{A} = \cup_{n=0}^{\infty} A_n = \mathcal{A}_1 * \mathcal{A}_2$, where $\mathcal{A}_1 * \mathcal{A}_2$ is the algebraic free-product of $\mathcal{A}_1$ and $\mathcal{A}_2$ with its evident involution.

Theorem 4.3. *Let (A_ι, L_ι) and $(\tilde{A}_\iota, \tilde{L}_\iota)$ be C^* -metric algebras coming from $(\{A_\iota\}, \sigma_\iota)$ and $(\{\tilde{A}_\iota\}, \tilde{\sigma}_\iota)$, respectively, with the faithful tracial states σ_ι and $\tilde{\sigma}_\iota$ and constant C for all $\iota = 1, 2$. Denote*

$$(A, \sigma) = (A_1, \sigma_1) * (A_2, \sigma_2),$$

$$(\tilde{A}, \tilde{\sigma}) = (\tilde{A}_1, \tilde{\sigma}_1) * (\tilde{A}_2, \tilde{\sigma}_2).$$

Suppose that $\alpha_1 : A_1 \rightarrow \tilde{A}_1$ and $\alpha_2 : A_2 \rightarrow \tilde{A}_2$ are two Lipschitz unital *-homomorphisms such that $\tilde{\sigma}_i \circ \alpha_i = \sigma_i$. Then there is a unique Lipschitz unital *-homomorphism $\alpha : A \rightarrow \tilde{A}$ such that for every $i \in \{1, 2\}$ the diagram

$$\begin{array}{ccccc} A_i & \xrightarrow{\lambda_i} & A & & \\ \alpha_i \downarrow & & \alpha \downarrow & \searrow \sigma & \\ \tilde{A}_i & \xrightarrow{\tilde{\lambda}_i} & \tilde{A} & \xrightarrow{\tilde{\sigma}} & \mathbb{C} \end{array}$$

commutes, where λ_i and $\tilde{\lambda}_i$ are the maps arising from the free product construction.

Proof. Consider the unital *-homomorphisms $\beta_1 = \tilde{\lambda}_1 \circ \alpha_1 : A_1 \rightarrow \tilde{A}$ and $\beta_2 = \tilde{\lambda}_2 \circ \alpha_2 : A_2 \rightarrow \tilde{A}$. We have the following diagrams commute:

$$\begin{array}{ccccc} & & \mathbb{C} & & \\ & \nearrow \sigma_1 & \uparrow \tilde{\sigma} & \nwarrow \sigma_2 & \\ A_1 & \xrightarrow{\beta_1} & \tilde{A} & \xleftarrow{\beta_2} & A_2 \end{array}$$

for $i \in \{1, 2\}$. By lemma 1.3 of [9], there is a *-homomorphism $\alpha : A \rightarrow \tilde{A}$ such that $\alpha \circ \lambda_i = \beta_i$ and $\tilde{\sigma} \circ \alpha = \sigma$. In particular, we have that $\alpha \circ \lambda_i = \tilde{\lambda}_i \circ \alpha_i$ and $\alpha(1_A) = \alpha(\lambda_1(1_{A_1})) = \beta_1(1_{A_1}) = (\tilde{\lambda}_1 \circ \alpha_1)(1_{A_1}) = 1_{\tilde{A}}$.

Since A is generated by $\lambda_1(A_1) \cup \lambda_2(A_2)$, it is clear that α will be unique if it exists. It is also clear that $\alpha(\mathcal{A}) \subseteq \tilde{\mathcal{A}}$, and so the Lipschitzness of α follows from Proposition 4.1, Proposition 4.2 and Theorem 1.1. $\square$

REFERENCES

1. E. M. Alfsen, *Compact convex sets and boundary integrals*, Springer-Verlag, Berlin, New York, 1971.
2. S. Baez, A. P. Balachandran, S. Vaidya, B. Ydri, *Monopoles and solitons in fuzzy physics*, Comm. Math. Phys., **208**(2000), no. 3, 787-798. arXiv:hep-th/9811169.
3. E. F. Blanchard, K. J. Dykema, *Embeddings of reduced free products of operator algebras*, Pacific J. Math., **199**(2001), no. 1, 1-19. arXiv:math.OA/9911012.
4. A. Connes, *Compact metric spaces, Fredholm modules, and hyperfiniteness*, Ergodic Theory Dynam. Systems, **9**(1989), no. 2, 207-220.
5. A. Connes, *Noncommutative geometry*, Academic Press, Inc., San Diego, CA, 1994.
6. A. Connes, *Noncommutative geometry and reality*, J. Math. Phys., **36**(1995), no. 11, 6194-6231.
7. J. B. Conway, *A course in functional analysis*, Springer-Verlag, New York, second edition, 1990.
8. R. S. Doran, V. A. Belfi, *Characterizations of C*-algebras. The Gelfand-Naimark theorems*, Monographs and Textbooks in Pure and Applied Mathematics, 101. Marcel Dekker, Inc., New York, 1986.
9. K. J. Dykema, M. Rørdam, *Projections in free product C*-algebras*, Geom. Funct. Anal., **8**(1998), 1-16. arXiv:funct-an/9702016.
10. Harald Grosse, Christian W. Rupp, Alexander Strohmaier, *Fuzzy line bundles, the Chern character and topological charges over the fuzzy sphere*, J. Geom. Phys., **42**(2002), no. 1-2, 54-63. arXiv:math-ph/0105033.
11. R. V. Kadison, *A representation theory for commutative topological algebra*, Mem. Amer. Math. Soc., **7**(1951).
12. L. V. Kantorovič, *On the translocation of masses*, C. R. (Doklady) Acad. Sci. URSS (N.S.), **37**(1942), 199-201.
13. L. V. Kantorovič, G. Š. Rubinštein, *On a functional space and certain extremum problems*, Dokl. Akad. Nauk SSSR (N.S.), **115**(1957), 1058-1061.
14. D. Kerr, *Matricial quantum Gromov-Hausdorff distance*, J. Funct. Anal., **205**(2003), no. 1, 132-167. arXiv:math.OA/0207282.

15. D. Kerr, *Dimension and dynamical entropy for metrized C^* -algebras*, Comm. Math. Phys., **232**(2003), no. 3, 501-534. arXiv:math.OA/0211043.
16. D. Kerr, H. Li, *On Gromov-Hausdorff convergence for operator metric spaces*, J. Operator Theory, to appear, arXiv:math.OA/0411157.
17. G. Landi, *An introduction to noncommutative spaces and their geometries*, Lecture Notes in Physics. New Series m: Monographs, 51. Springer-Verlag, Berlin, 1997. arXiv:hep-th/9701078.
18. H. Li, *Order-unit quantum Gromov-Hausdorff distance*, J. Funct. Anal., **231**(2006), no. 2, 312-360. arXiv:math.OA/0312001.
19. H. Li, *C^* -algebraic quantum Gromov-Hausdorff distance*, arXiv:math.OA/0312003.
20. H. Li, *Compact quantum metric spaces and ergodic actions of compact quantum groups*, J. Funct. Anal., to appear, arXiv:math.OA/0411178.
21. H. Li, *Metric aspects of noncommutative homogeneous spaces*, arXiv:0810.4694.
22. N. Ozawa, M. A. Rieffel, *Hyperbolic group C^* -algebras and free-product C^* -algebras as compact quantum metric spaces*, Canad. J. Math., **57**(2005), no. 5, 1056-1079. arXiv:math.OA/0302310.
23. V. I. Paulsen, *Completely bounded maps and dilations*, Pitman Research Notes in Mathematics Series, 146. Longman Scientific & Technical, Harlow; John Wiley & Sons, Inc., New York, 1986.
24. M. A. Rieffel, *Metrics on states from actions of compact groups*, Doc. Math., **3**(1998), 215-229. arXiv:math.OA/9807084.
25. M. A. Rieffel, *Metrics on state spaces*, Doc. Math., **4**(1999), 559-600. arXiv:math.OA/9906151.
26. M. A. Rieffel, *Group C^* -algebras as compact quantum metric spaces*, Doc. Math., **7**(2002), 605-651. arXiv:math.OA/0205195.
27. M. A. Rieffel, *Gromov-Hausdorff distance for quantum metric spaces*, Mem. Amer. Math. Soc. **168**(2004), no. 796, 1-65. arXiv:math.OA/0011063.
28. M. A. Rieffel, *Matrix algebras converge to the sphere for quantum Gromov-Hausdorff distance*, Mem. Amer. Math. Soc. **168**(2004), no. 796, 67-91. arXiv:math.OA/0108005.
29. M. A. Rieffel, *compact quantum metric spaces*, Operator algebras, quantization, and noncommutative geometry, 315-330, Contemp. Math., 365, Amer. Math. Soc., Providence, RI, 2004. arXiv:math.OA/0308207.
30. M. A. Rieffel, *Vector bundles and Gromov-Hausdorff distance*, arXiv:math.MG/0608266.
31. M. A. Rieffel, *Leibniz seminorms for "Matrix algebras converge to the sphere"*, arXiv:0707.3229.
32. P. Valtancoli, *Projectors for the fuzzy sphere*, Modern Phys. Lett. A, **16**(2001), no. 10, 639-645. arXiv:hep-th/0101189.
33. J. C. Várilly, *An introduction to noncommutative geometry*, EMS Series of Lectures in Mathematics. European Mathematical Society (EMS), Zürich, 2006. arXiv:physics/9709045.
34. D. V. Voiculescu, *Symmetries of some reduced free product C^* -algebras*, Operator algebras and their connections with topology and ergodic theory (Buşteni, 1983), 556-588, Lecture Notes in Math., 1132, Springer, Berlin, 1985.
35. D. V. Voiculescu, *On the existence of quascentral approximate units relative to normed ideals. I*, J. Funct. Anal., **91**(1990), no. 1, 1-36.
36. D. V. Voiculescu, K. J. Dykema, A. Nica, *Free random variables*, A noncommutative probability approach to free products with applications to random matrices, operator algebras and harmonic analysis on free groups. CRM Monograph Series, 1. American Mathematical Society, Providence, RI, 1992.
37. W. Wu, *Locally pre- C^* -equivalent algebras*, Proc. Amer. Math. Soc., **131**(2003), no. 2, 555-562.
38. W. Wu, *Non-commutative metric topology on matrix state spaces*, Proc. Amer. Math. Soc. **134**(2006), no. 2, 443-453. arXiv:math.OA/0410587.
39. W. Wu, *Quantized Gromov-Hausdorff distance*, J. Funct. Anal. **238**(2006), no. 1, 58-98. arXiv:math.OA/0503344.

DEPARTMENT OF MATHEMATICS, EAST CHINA NORMAL UNIVERSITY, SHANGHAI 200241, P.R. CHINA

E-mail address: wwu@math.ecnu.edu.cn